


Breeding durable eucalypts

Luis Apiolaza, Clemens Altaner, Jackley Li & Nick Davies

School of Forestry | Kura Ngahere

University of Canterbury, Christchurch

New Zealand | Aotearoa



Where are we?

Madrid

Barcelona

Montpellier

Woodville trial

Lawson's trial



Motivation





canadulensis

tricarpa

quadrangulata

macrorrhyncha

canadulensis

tricarpa

globoides

argophylla

canadulensis

cladocalyx

eugenoides

boissipana

quadrangulata

macrorrhyncha

eugenoides

longifolia

Demonstration trial
(photo: Rick Alexander)

It can get fairly dry at times...



How is our breeding programme structured (for **each of 5 species**)?

Breeding Trials
(3+ sites)
Like Lawson's



Selection of
best trees
from all
families

Growth-strain nursery trial
(1 site: Woodville)



Family-level
screening

Selections

Based on individual
tree assessment
+
family-level
assessments



Lawson's Progeny Trial (*E. bosistoana*)

Durability (using 3 criteria)



Heartwood variability ➤ Durability variability

Heartwood
presence/absence


Heartwood quantity

Heartwood quality
(extractives content
via NIR)

Heartwood quality
(mass loss for brown
& white rot via NIR)




Environmental effects on heartwood presence/absence




Heritability for heartwood presence ~ 0.2 — 0.3

Heartwood quantity (heart diameter)



Heritability for heartwood diameter ~ 0.4 — 0.5

Relationship between the durability criteria



Growth strain: nursery trial





Larger opening ➤ Higher growth strain




Chauhan & Entwistle. 2010. Measurement of surface growth stress in *Eucalyptus nitens* Maiden by splitting a log along its axis. *Holzforschung*, 64(2).

DOI:10.1515/hf.2010.022

Davies, Apiolaza & Sharma. 2017 Heritability of growth strain in *Eucalyptus bosistoana*: a Bayesian approach with left-censored data. *New Zealand Journal of Forestry Science* 47:5 DOI: 10.1186/s40490-017-0086-

Growth strain: some results




Growth strain heritability ~ 0.3

Our most disappointing trait: right now the resolution is only enough to discriminate between best and worst families

In summary

- We are domesticating 5 species, focusing on the quality of solid-products.
- We assess early 2 to 7 years.
- We match trees to environments: niche products for niche sites.
- Durability split into three traits: 1/0, heartwood diameter and extractives content.
- Growth strain is much harder to deal with; right now can tell between best & worst families, but not at the tree level.





With thanks to Monika Sharma, Ruth McConnochie, Paul Millen
& Shaf van Ballekom

SUPPORTERS:



Ministry for Primary Industries
Manatū Ahu Matua

